

The lot number #LVSDG-0100 is a lentivirus-based pseudovirus pseudotyped with the SARS-CoV-2 D614G variant spike protein. This quality control report demonstrates that the lot #LVSDG-0100 is efficient for cell transduction and can be effectively neutralized by a standard neutralizing antibody.

1. Transduction efficiency assay

Target cells	HEK293 cells (ACE2 ⁺ , TMPRSS2 ⁺)
Volume of pseudoviruses	5 µL/well
Detection signal	Fluorescence (GFP)
Detection method	Fluorescent microscopy





Figure 1: GFP-positive cells after transduction with the LV-Pseudovirus #LVSDG-0100.

A volume of 5 µL of LV-pseudovirus was mixed to 45 µL of culture complete medium, in a 96-well plate. Then, an additional volume of 50 µL containing 30 000 HEK293 cells (ACE2⁺, TMPRSS2⁺) were seeded in each well. GFP expression analysis was performed 48 hours post-infection by fluorescent microscopy.

Conclusion

The production batch of LV-Pseudoviruses #LVSDG-0100 can transduce the target cells.

2. Neutralization assay

Target cells	HEK293 cells (ACE2 ⁺ , TMPRSS2 ⁺)
Volume of pseudoviruses	5 µL/well
Neutralizing antibody standard	Anti-Spike Protein (RBD) [CV30], AB02019-12.1
Detection signal	Fluorescence (GFP)
Detection method	Flow cytometry




Figure 2 : Neutralization curve of the batch #LVSDG-0100.

A neutralizing antibody ([AB02019-12.1](#)) was serially diluted in a final volume of 50 µL of complete medium and incubated for 1 hour with 5 µL of LV-pseudoviruses, in a 96-well plate. Then, an additional 50 µL containing 30 000 HEK293 cells (ACE2⁺, TMPRSS2⁺) were seeded in each well and incubated for 48 hours before flow cytometry analysis. Data in relative unit fluorescence (RFU) were obtained from the analysis of 10 000 cells. Control conditions with pseudoviruses or cells only were also included in the assay as negative controls. The normalized values are analyzed by choosing a nonlinear regression analysis followed by a log (inhibitor) vs response – variable slope function in GraphPad Prism software. Data are representative of duplicates.

Conclusion

The production batch of LV-Pseudoviruses #LVSDG-0100 can be efficiently neutralized by a neutralizing antibody.

3. Additional information

Caution

We recommend determining the optimal pseudovirus volume to use according to your specific experimental conditions.

LV-pseudovirus

3rd generation, replication incompetent

Pseudotyping

SARS-CoV-2 D614G variant spike protein (GenBank: [MN908947](#)) with the D614G mutation. The spike protein has an 18-aa cytoplasmic tail truncation for optimal infection.

Glycosylation origin

Human

Reporter protein

Green fluorescent protein

Contact us for more information mathias.mangion@ivanobioscience.com